

Constraints on Gluon Sivers Distribution from RHIC Results¹

M. Anselmino*, U. D'Alesio[†], S. Melis[†] and F. Murgia[†]

**Dipartimento di Fisica Teorica, Università di Torino and
INFN, Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy*

*[†]Dipartimento di Fisica, Università di Cagliari and
INFN, Sezione di Cagliari, C.P. 170, I-09042 Monserrato (CA), Italy*

Abstract. We consider the recent RHIC data on the transverse single spin asymmetry (SSA) A_N , measured in $p^\uparrow p \rightarrow \pi^0 X$ processes at mid-rapidity by the PHENIX Collaboration. We analyze this experimental information within a hard scattering approach based on a generalized QCD factorization scheme, with unintegrated, transverse momentum dependent (TMD), parton distribution and fragmentation functions. In this kinematical region, only the gluon Sivers effect could give a large contribution to A_N ; its vanishing value is thus used to give approximate upper limits on the gluon Sivers function (GSF). Additional constraints from the Burkardt sum rule for the Sivers distributions are also discussed.

Keywords: single spin asymmetries, TMD distributions, Sivers effect

PACS: 12.38.Bx, 13.88.+e, 13.85.Ni, 14.70.Dj

FORMAL APPROACH

Transverse single spin asymmetries can originate, even with a short distance helicity conserving pQCD dynamics, from spin- $\mathbf{k}_\perp$ correlations in the soft components of the hadronic process $A^\uparrow B \rightarrow CX$. According to the hard scattering approach to hadronic interactions developed in Refs. [1, 2, 3], based on the assumption of a generalized QCD factorization scheme which involves unintegrated TMD parton distribution and fragmentation functions, the general structure of the cross section for the polarized hadronic process $(A, S_A) + (B, S_B) \rightarrow C + X$ reads (see Ref. [3] for more details)

$$\begin{aligned} \frac{E_C d\sigma^{(A,S_A)+(B,S_B) \rightarrow C+X}}{d^3\mathbf{p}_C} &= \sum_{a,b,c,d,\{\lambda\}} \int \frac{dx_a dx_b dz}{16\pi^2 x_a x_b z^2 s} d^2\mathbf{k}_{\perp a} d^2\mathbf{k}_{\perp b} d^3\mathbf{k}_{\perp C} \delta(\mathbf{k}_{\perp C} \cdot \hat{\mathbf{p}}_C) \\ &\times J(k_{\perp C}) \rho_{\lambda_a, \lambda'_a}^{a/A, S_A} \hat{f}_{a/A, S_A}(x_a, \mathbf{k}_{\perp a}) \rho_{\lambda_b, \lambda'_b}^{b/B, S_B} \hat{f}_{b/B, S_B}(x_b, \mathbf{k}_{\perp b}) \\ &\times \hat{M}_{\lambda_c, \lambda_d; \lambda_a, \lambda_b} \hat{M}_{\lambda'_c, \lambda'_d; \lambda'_a, \lambda'_b}^* \delta(\hat{s} + \hat{t} + \hat{u}) \hat{D}_{\lambda_c, \lambda'_c}^{\lambda_c, \lambda'_c}(z, \mathbf{k}_{\perp C}), \quad (1) \end{aligned}$$

where all parton intrinsic motions are fully taken into account, both in the soft, non perturbative components and in the hard, pQCD interactions.

The main features of Eq. (1) are the appearance of several spin and $\mathbf{k}_\perp$ dependent distribution and fragmentation functions and the non-collinear partonic configuration

¹ Talk delivered by U. D'Alesio at the "17th International Spin Physics Symposium", SPIN2006, October 2-7, 2006, Kyoto, Japan.

which lead to many $\mathbf{k}_\perp$ dependent phases. In Ref. [3] it was explicitly shown that the only sizeable contributions to the transverse single spin asymmetry $A_N(p^\uparrow p \rightarrow \pi X)$, in the kinematical region of large positive x_F come from the Siverts [4] and, less importantly, from the Collins [5] mechanisms. Moreover, while the quark contribution is totally dominant at large, positive x_F values (for polarized protons moving along the positive Z-axis), the gluon contribution may be sizeable in the mid-rapidity and negative x_F regions.

Data in the mid-rapidity region are available from the E704 [6] and PHENIX [7] experiments; the region of negative values of x_F has been covered by the STAR [8] and BRAHMS experiments [9]. In these kinematical regions $A_N(p^\uparrow p \rightarrow \pi X)$ is largely dominated by the Siverts effect [4], all other contributions being almost vanishing, and Eq. (1) gives [1]:

$$\begin{aligned} \frac{E_\pi d\sigma^\uparrow}{d^3\mathbf{p}_\pi} - \frac{E_\pi d\sigma^\downarrow}{d^3\mathbf{p}_\pi} \simeq \sum_{a,b,c,d} \int \frac{dx_a dx_b dz}{\pi x_a x_b z^2 s} d^2\mathbf{k}_{\perp a} d^2\mathbf{k}_{\perp b} d^3\mathbf{k}_{\perp \pi} \delta(\mathbf{k}_{\perp \pi} \cdot \hat{\mathbf{p}}_c) J(k_{\perp \pi}) \\ \times \Delta \hat{f}_{a/p^\uparrow}(x_a, \mathbf{k}_{\perp a}) \hat{f}_{b/p}(x_b, k_{\perp b}) \hat{s}^2 \frac{d\hat{\sigma}^{ab \rightarrow cd}}{d\hat{t}}(x_a, x_b, \hat{s}, \hat{t}, \hat{u}) \delta(\hat{s} + \hat{t} + \hat{u}) \hat{D}_{\pi/c}(z, k_{\perp \pi}), \end{aligned} \quad (2)$$

where

$$\Delta \hat{f}_{a/p^\uparrow}(x_a, \mathbf{k}_{\perp a}) \equiv \hat{f}_{a/p^\uparrow}(x_a, \mathbf{k}_{\perp a}) - \hat{f}_{a/p^\downarrow}(x_a, \mathbf{k}_{\perp a}) = \Delta^N \hat{f}_{a/p^\uparrow}(x_a, k_{\perp a}) \cos \phi_a. \quad (3)$$

$\Delta^N \hat{f}_{a/p^\uparrow}(x_a, k_{\perp a})$ [or $f_{1T}^\perp(x_a, k_{\perp a})$] is referred to as the Siverts distribution function of parton a inside a transversely polarized (along the Y-axis) proton (moving along the Z-axis). ϕ_a is the azimuthal angle of the intrinsic transverse momentum $\mathbf{k}_{\perp a}$ of parton a .

The azimuthal phase factor $\cos \phi_a$ appearing in the numerator of A_N , Eqs. (2) and (3), plays a crucial role and deserves a comment. The only other term depending on ϕ_a in Eq. (2) is the partonic cross section, in particular via the corresponding Mandelstam variable $\hat{t}$. Therefore while at large positive x_F (t -channel dominated) the integration over ϕ_a does not necessarily suppress A_N , for negative values of x_F (u -channel dominated) one is roughly left with the $d^2\mathbf{k}_{\perp a} \cos \phi_a$ integration alone, which cancels the potentially large Siverts contribution. As a consequence, one cannot get significant information on the gluon Siverts distribution from the recent STAR and BRAHMS data at negative values of x_F . (Notice that at lower values of $\sqrt{s}$ the suppression induced by the $\cos \phi_a$ dependence would be less drastic.)

The same arguments do not apply to inclusive hadronic processes at mid-rapidity and moderately large p_T values, for which data from PHENIX [7] are already available, for neutral pions and charged hadron production. For these processes, the gluon contribution is dominant and the Siverts effect can survive the phase integration. The possibility of accessing the gluon Siverts function has been also investigated in Refs. [10, 11, 12].

Indirect constraints on the GSF could also be obtained from a sum rule for the Siverts distribution recently derived by Burkardt [13]. The Burkardt Sum Rule (BSR) states that the total (integrated over x and $\mathbf{k}_\perp$) transverse momentum of all partons (quarks, antiquarks and gluons) in a transversely polarized proton must be zero,

$$\langle \mathbf{k}_\perp \rangle = \sum_a \langle \mathbf{k}_\perp \rangle_a = \int dx \int d^2\mathbf{k}_\perp \mathbf{k}_\perp \sum_a \Delta \hat{f}_{a/p^\uparrow}(x, \mathbf{k}_\perp) = 0. \quad (4)$$

In the following we shall simply check whether or not the proposed parameterizations of the Siverson functions are compatible with the BSR.

PHENOMENOLOGY

We consider the PHENIX data [7] on A_N for the $p^\uparrow p \rightarrow \pi^0 X$ process at RHIC, at $\sqrt{s} = 200$ GeV, with p_T ranging from 1.0 to 5.0 GeV/c and mid-rapidity values, $|\eta| < 0.35$. In this kinematical regime, at the lowest p_T values, $x_a^{\min}$ can be as small as 0.005. Therefore, partonic channels involving a gluon in the transversely polarized initial proton dominate over those involving a quark. This gluon dominance, together with the (almost) vanishing of all possible contributions to A_N other than the Siverson effect, allows to put upper bounds on the GSF. Due to possible mixing with quark initiated contributions, the same is not true for the E704 data [6] at lower energies and comparable rapidity and p_T ranges.

In Ref. [1] we have shown that reasonable fits to the SSA for the $p^\uparrow p \rightarrow \pi X$ process at large positive x_F can be obtained by using valence-like Siverson functions for u and d quarks, which turn out to have opposite signs. The use of valence-like u and d Siverson functions alone predicts an almost vanishing SSA in the mid-rapidity region, compatible with available data (see Fig. 1(left), for the PHENIX results). Their parameterizations [1, 14] are also compatible with the BSR, Eq. (4). Although a large gluon Siverson function would not modify the analysis of the SIDIS, E704 and STAR data at large positive x_F , it would strongly affect the description of the mid-rapidity PHENIX data.

In what follows (see Ref. [15] for details) we therefore try to understand what is the maximum value of $|\Delta^N \hat{f}_{g/p^\uparrow}(x, k_\perp)|/2\hat{f}_{g/p}(x, k_\perp)$ allowed by the PHENIX data; our results are summarized in Fig. 1 (left panel for the SSA and right panel for the GSF):

- The thin, solid line in Fig. 1(left), results from computing A_N using only the valence-like u and d Siverson functions of Ref. [1].
- The dot-dashed curve in Fig. 1(left) has been obtained by saturating (in magnitude) the GSF to the natural positivity bound [see Eq. (3)]

$$\Delta^N \hat{f}_{g/p^\uparrow}(x, k_\perp) = -2\hat{f}_{g/p}(x, k_\perp). \quad (5)$$

The sea-quark Siverson functions are again assumed to vanish. This choice leads to a SSA definitely in contradiction with data and to a strong violation of the BSR.

- The thick, solid curve in Fig. 1(left) has been obtained still assuming that there is no sea-quark Siverson contribution, and looking for a parameterization of $\Delta^N \hat{f}_{g/p^\uparrow}$ yielding values of A_N falling, approximately, within one-sigma deviation below the lowest p_T data. The corresponding x -dependent part of the GSF, normalized to its positivity bound, $|\Delta^N \hat{f}_{g/p^\uparrow}(x)|/2\hat{f}_{g/p}(x)$, is shown as the solid curve in Fig. 1(right). It leads, within the x range covered by the data, to a strong violation of the BSR.
- We finally consider the inclusion of all sea-quarks ($u_s, \bar{u}, d_s, \bar{d}, s, \bar{s}$) contributions by using a non-vanishing positive Siverson function which saturates the positivity bound [that is, $\Delta^N \hat{f}_{q_s/p^\uparrow}(x, k_\perp) \equiv 2\hat{f}_{q_s/p}(x, k_\perp)$]. These contributions could then cancel the negative contribution to A_N of a possibly large GSF. We then look for the largest negative GSF which, together with a positive maximized sea-quark contribution, leads again to the




FIGURE 1. Left: the computed SSA, A_N , compared with PHENIX data [7], with different choices for the gluon and sea-quark Siversons functions. Right: the value of the normalized GSF, $|\Delta^N f_{g/p^+}(x)|/2f_{g/p}(x)$.

SSA represented by the thick, solid line of Fig. 1(left). This curve results now as the sum of the (maximized) sea and valence quark contribution (dotted curve) and that of the new GSF (dashed curve), which is plotted as the dashed curve in Fig. 1(right). It is the largest (overmaximized) gluon Siversons function compatible with PHENIX data. Within the x range covered by the data, the (over)maximized sea-quark Siversons distributions give a positive contribution which strongly suppresses the negative contribution of the GSF, so that in this scenario the BSR is satisfied within a 10% level of accuracy.

Summarizing, our analysis shows that the PHENIX data on $A_N(p^\uparrow p \rightarrow \pi^0 X)$ allow to put significant quantitative bounds on the magnitude of the GSF. Similar conclusions have been recently reached by studying the Siversons effect in SIDIS off a deuteron target [16].

REFERENCES

1. U. D'Alesio and F. Murgia, *Phys. Rev.* **D70**, 074009 (2004).
2. M. Anselmino, M. Boglione, U. D'Alesio, E. Leader and F. Murgia, *Phys. Rev.* **D71**, 014002 (2005).
3. M. Anselmino, M. Boglione, U. D'Alesio, E. Leader, S. Melis and F. Murgia, *Phys. Rev.* **D73**, 014020 (2006).
4. D.W. Siversons, *Phys. Rev.* **D41**, 83 (1990); **D43**, 261 (1991).
5. J.C. Collins, *Nucl. Phys.* **B396**, 61 (1993).
6. D.L. Adams *et al.* (FNAL-E704 Collaboration), *Phys. Rev.* **D53**, 4747 (1996).
7. S.S. Adler *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **95**, 202001 (2005).
8. C.A. Gagliardi (for the STAR Collaboration), hep-ex/0607003.
9. F. Videbaek (for the BRAHMS Collaboration), nucl-ex/0601008.
10. M. Anselmino, M. Boglione, U. D'Alesio, E. Leader and F. Murgia, *Phys. Rev.* **D70**, 074025 (2004).
11. D. Boer and W. Vogelsang, *Phys. Rev.* **D69**, 094025 (2004).
12. I. Schmidt, J. Soffer and J.J. Yang, *Phys. Lett.* **B612**, 258 (2005).
13. M. Burkardt, *Phys. Rev.* **D69**, 057501 (2004); **D69**, 091501 (2004).
14. M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia and A. Prokudin, *Phys. Rev.* **D71**, 074006 (2005); **D72**, 094007 (2005).
15. M. Anselmino, U. D'Alesio, S. Melis and F. Murgia, *Phys. Rev.* **D74**, 094011 (2006).
16. S.J. Brodsky and S. Gardner, hep-ph/0608219.